

YOUR GUIDE TO TREATMENT

# STEP BY STEP

Please see Important Safety Information on pages 11-13 and full [Prescribing Information/Medication Guide](#), including serious side effects.

  
**LEMTRADA**<sup>®</sup>  
alemtuzumab <sup>12mg</sup> iv

BEFORE TREATMENT

DURING TREATMENT

AFTER TREATMENT

FAQS AND IMPORTANT  
SAFETY INFORMATION

## PLANNING FOR

**WHAT'S NEXT**

Reading the information in this guide will help with your preparation before, during, and after your LEMTRADA treatment.

**INDICATION**

LEMTRADA is a prescription medicine used to treat relapsing forms of multiple sclerosis (MS), to include relapsing-remitting disease and active secondary progressive disease, in adults. Since treatment with LEMTRADA can increase your risk of getting certain conditions and diseases, LEMTRADA is generally prescribed for people who have tried 2 or more MS medicines that have not worked well enough. LEMTRADA is not recommended for use in patients with clinically isolated syndrome (CIS). It is not known if LEMTRADA is safe and effective for use in children under 17 years of age.

**SELECTED IMPORTANT SAFETY INFORMATION**

LEMTRADA can cause serious side effects including autoimmune problems, infusion reactions, stroke, tears in your arteries that supply blood to your brain (carotid and vertebral arteries), some kinds of cancers, thyroid problems, low blood counts (cytopenias), inflammation of the liver, hemophagocytic lymphohistiocytosis, serious infections, progressive multifocal leukoencephalopathy (PML), inflammation of the gallbladder without gallstones (acalculous cholecystitis), and swelling of lung tissue (pneumonitis). Because of the risks of autoimmune problems, infusion reactions and some kinds of cancers, LEMTRADA is only available through a restricted program called the Risk Evaluation and Mitigation Strategy (REMS) Program.

**WHAT'S DONE**

**DETERMINE WITH HCP IF LEMTRADA IS RIGHT FOR ME**



**ENROLL IN LEMTRADA REMS**



**READY TO BEGIN TAKING LEMTRADA**

Please see Important Safety Information on pages 11-13 and full [Prescribing Information/Medication Guide](#), including serious side effects.

**LEMTRADA**<sup>®</sup>  
alemtuzumab<sup>12mg</sup> iv

# EVERY STEP OF THE WAY THERE'S SUPPORT

It's comforting to have someone to talk to. Your *MS One to One*® nurse will be available, after you sign the LEMTRADA Services Form, to help you with:

- ✓ Verifying insurance benefits and answering other questions
- ✓ Locating an infusion center
- ✓ Organizing and preparing for your treatment
- ✓ Explaining what to expect after your infusion days
- ✓ Setting reminders for your monthly monitoring

## FINANCIAL SUPPORT

To help make treatment costs and co-pays more affordable, the **LEMTRADA Co-Pay Program offers financial assistance to help eligible,\* commercially insured patients receive:**

**\$0** OUT-OF-POCKET COST  
FOR LEMTRADA

**\$100** PER DAY FOR CHARGES  
RELATED TO THE  
ADMINISTRATION OF  
YOUR LEMTRADA INFUSION  
UP TO

\*Patients who have coverage or prescriptions paid for in part or in full under Medicare, Medicaid, or other state or federally funded healthcare are not eligible. The LEMTRADA Co-Pay Program is subject to termination or modification at any time. If you are not eligible for the LEMTRADA Co-Pay Program and need help with out-of-pocket expenses, *MS One to One* can help review your coverage options. Depending on your specific situation, your *MS One to One* nurse can direct you to other patient assistance programs that may offer you financial support. Treatment related, infusion out-of-pocket costs are not reimbursable in MA, MI, MN, or RI.

Please see Important Safety Information on pages 11-13 and full [Prescribing Information/Medication Guide](#), including serious side effects.

# MS ONE TO ONE® NURSES ARE AVAILABLE 24/7

*MS One to One* will provide you with access to a nurse anytime. Regular call center hours are Mon-Fri, 8:30 am-8:00 pm ET at **1-855-676-6326**.



name

phone number



name

address

phone number



Even if your dedicated *MS One to One* nurse is not available, someone is always there to answer questions.

**LEMTRADA**  
alemtuzumab<sup>12mg</sup>

# JOURNEY TO TREATMENT CHECKLIST

## 6 WEEKS BEFORE



### SCHEDULE APPOINTMENTS

Schedule your LEMTRADA infusion appointments. Ask about policies on bringing food, entertainment, visitors, etc.



### CALL YOUR NURSE

Call *MS One to One*® at 1-855-676-6326 and give your nurse your appointment dates



### GET PRETREATMENT TESTS AND VACCINATIONS

Complete blood and urine tests and a skin check as prescribed. Get any other tests and vaccinations your healthcare team has recommended



### MAKE PLANS

Arrange for transportation, work coverage, and/or child care for days of infusion

## 1 WEEK BEFORE



### CONFIRM DATES

Confirm your infusion appointments with your infusion center



### GET MEDICATIONS

Get medications as prescribed by your healthcare team



### CONFIRM PLANS

Confirm transportation to and from your infusion appointments. Do not drive yourself

## 1 DAY BEFORE



### PACK

- headphones
- book
- food/snacks
- bottled water or refillable container



### DOWNLOAD

- movies
- podcasts
- music playlist

### MEDICATIONS YOUR DOCTOR WILL PRESCRIBE INCLUDE:

**Corticosteroids** to lower your chances of getting serious infusion reactions, your healthcare provider will give you a medicine called corticosteroids before your first 3 infusions of a treatment course. **Antihistamines and fever reducers** may be given prior to your infusion to reduce the chances of infusion reactions, or to treat them if they happen. **Antivirals** will be prescribed to you to take starting on your first day of infusions. Continue taking medications as recommended by your healthcare team.

Please see Important Safety Information on pages 11-13 and full [Prescribing Information/Medication Guide](#), including serious side effects.

**LEMTRADA**<sup>®</sup>  
alemtuzumab<sup>12mg</sup><sub>iv</sub>

TODAY IS

# TREATMENT DAY

You've done your research, spoken up to advocate for your treatment goals, and made all necessary arrangements.

Relax and remember...

YOU'VE GOT  
**THIS!**

## TREATMENT CHECKLIST

 **PACK BAG** **WEAR COMFY  
CLOTHES** **TAKE RECOMMENDED  
PRE-MEDICATIONS** **ARRIVE EARLY** **STAY HYDRATED**

Please see Important Safety Information on pages 11-13 and full [Prescribing Information/Medication Guide](#), including serious side effects.

**LEMTRADA**<sup>®</sup>  
alemtuzumab<sup>12mg</sup> iv

# STEPS TO HELP MANAGE INFUSION REACTIONS

LEMTRADA can cause serious infusion reactions that may cause death. Serious infusion reactions may happen while you receive, or up to 24 hours or longer after you receive LEMTRADA.

- **92%** of patients experienced infusion reactions in the LEMTRADA clinical trials
- **3%** of these reactions were serious



## MANAGE INFUSION REACTIONS

You will receive your infusion at a healthcare facility with equipment and staff trained to manage infusion reactions, including serious allergic reactions, and urgent heart or breathing problems.



## MONITORING INFUSION REACTIONS

You will be watched while you receive, and for 2 hours or longer after you receive, LEMTRADA on each infusion day. If a serious infusion reaction happens while you are receiving LEMTRADA, your infusion may be stopped.



## REMINDER

**Make sure you stay hydrated before, during, and after your infusion.**

Please see Important Safety Information on pages 11-13 and full [Prescribing Information/Medication Guide](#), including serious side effects.

# WATCH FOR THESE SYMPTOMS

Tell your healthcare team right away if you have any of the following symptoms of a serious infusion reaction during or after treatment:



**SWELLING IN YOUR MOUTH OR THROAT**



**TROUBLE BREATHING**



**WEAKNESS**



**FAST, SLOW, OR IRREGULAR HEARTBEAT**



**CHEST PAIN**



**RASH**

Some people have had serious and sometimes deadly strokes and tears in their carotid or vertebral arteries within 3 days of receiving LEMTRADA. Get help right away if you have any of the following symptoms that may be signs of a stroke or tears in your carotid or vertebral arteries: drooping of parts of your face, weakness on one side, sudden severe headache, difficulty with speech or neck pain.

**LEMTRADA**<sup>®</sup>  
alemtuzumab<sup>12mg</sup><sub>iv</sub>

# YOUR TREATMENT COURSE IS **COMPLETE!**

Self-care is top priority in the days and months following your infusion.

Take the time to...

## POST-TREATMENT TO-DO

- ✓ **REST, RELAX,  
AND TAKE CARE  
OF YOURSELF**
- ✓ **SCHEDULE  
MONITORING  
APPOINTMENTS**
- ✓ **PERFORM SYMPTOM  
SELF-CHECKS**



Please see Important Safety Information on pages 11-13 and full [Prescribing Information/Medication Guide](#), including serious side effects.

**LEMTRADA**<sup>®</sup>  
alemtuzumab<sup>12mg</sup> iv

# CARE FOR YOURSELF AFTER YOUR INFUSIONS

Be sure to follow any post-treatment instructions from your healthcare team, which may include:

- ✓ Washing hands frequently to reduce risk of infections
- ✓ Staying hydrated
- ✓ Resting
- ✓ Eating regularly
- ✓ Avoiding or adequately heating foods that may contain listeria during and after LEMTRADA treatment\*

## SELECTED IMPORTANT SAFETY INFORMATION

LEMTRADA can cause serious side effects including autoimmune problems, infusion reactions, stroke, tears in your arteries that supply blood to your brain (carotid and vertebral arteries), some kinds of cancers, thyroid problems, low blood counts (cytopenias), inflammation of the liver, hemophagocytic lymphohistiocytosis, serious infections, progressive multifocal leukoencephalopathy (PML), inflammation of the gallbladder without gallstones (acalculous cholecystitis), and swelling of lung tissue (pneumonitis). Because of the risks of autoimmune problems, infusion reactions and some kinds of cancers, LEMTRADA is only available through a restricted program called the Risk Evaluation and Mitigation Strategy (REMS) Program.

\*Listeria infections have occurred as early as 3 days after treatment and up to 8 months after the last LEMTRADA dose. For more information, please see page 10.

Please see Important Safety Information on pages 11-13 and full [Prescribing Information/Medication Guide](#), including serious side effects.

# TAKE AN ACTIVE ROLE IN COMPLETING YOUR MONITORING

Certain side effects of LEMTRADA can happen months and even years after treatment. That's why it is important to begin monitoring after your first infusion and continue regular monitoring until 4 years after your last infusion. Monitoring can help your healthcare team find potential side effects early.



Monthly blood and urine tests



Ongoing symptom self-checks



A yearly skin examination

## MANY OPTIONS FOR MONTHLY MONITORING

*MS One to One*<sup>®</sup> offers a range of options for monthly blood and urine testing at no cost to you.



**HOME:** Lab tech visits in the comfort of your home or office



**LAB:** Any local LabCorp<sup>®</sup> or Quest Diagnostics<sup>®</sup> patient service center\*



**HEALTHCARE PROVIDER'S OFFICE:** Samples can be taken at your healthcare team's office. With this option, you may incur a co-pay for your doctor's office visit. However, the cost of analyzing the samples will be covered by Sanofi Genzyme.

\*Trademarks not owned by Sanofi Genzyme corporation are the property of their respective owners.

**LEMTRADA**<sup>®</sup>  
alemtuzumab<sup>12mg</sup> iv



## IDENTIFY SYMPTOMS

# WITH SELF-CHECKS

Spot any physical changes by performing regular body checks at home by yourself in a mirror, or with the help of a family member or friend.

### EYES, NOSE, AND MOUTH

- Yellowing of the whites of your eyes
- Eye issues including swelling and/or vision problems
- Bleeding from gums or nose that is new or takes longer than usual to stop
- Swelling in your mouth or throat
- Drooping of parts of your face

### PAIN

- Neck pain
- Chest pain or tightness
- Sudden severe headache
- Stomach pain or discomfort

### BOWEL AND URINARY ISSUES

- Constipation
- Blood in the urine (red or tea-colored)
- Decrease in urine
- Dark urine

### SKIN


- Easy bruising
- Small, scattered spots that are red, pink, or purple
- Yellowing of the skin
- Bleeding from a cut that is hard to stop
- Swelling in your legs or feet
- Swelling in your neck
- New lump
- Rash

Please see Important Safety Information on pages 11-13 and full [Prescribing Information/Medication Guide](#), including serious side effects.



## REMINDER

Contact your healthcare team if you experience any of these symptoms, or new or worsening medical problems that have lasted several days, including changes in energy or mood, problems with thinking and/or balance.



### RESPIRATORY AND THROAT

- Cough that is not caused by a cold
- Coughing up blood
- Trouble swallowing or breathing
- Hoarseness or other voice changes that don't go away
- Swelling in your throat
- Wheezing
- Shortness of breath

### OTHER

- Fast, slow, or irregular heartbeat
- Excessive sweating or feeling cold
- Weakness
- Problems using arms or legs
- Worsening tiredness
- Nervousness
- Unexplained weight loss or gain
- Loss of appetite
- Fever
- Swollen glands
- For women: heavier menstrual periods than normal
- Nausea or vomiting
- Weakness on one side
- Difficulty with speech
- Fatigue

# WE HAVE ANSWERS

## HOW LONG DOES EACH INFUSION LAST?

Each infusion takes approximately 4 hours, and your healthcare team will monitor you for at least 2 hours for infusion reactions.

## WHAT KINDS OF SERIOUS SIDE EFFECTS CAN HAPPEN DURING AN INFUSION?

Tell your healthcare provider right away if you have any of the following symptoms of a serious infusion reaction during the infusion, and after you have left the healthcare facility: swelling in your mouth or throat, trouble breathing, weakness, fast, slow or irregular heartbeat, chest pain or rash.

## IS THERE ANYTHING I CAN DO TO HELP ADDRESS INFUSION SIDE EFFECTS?

- Your healthcare provider will give you a medicine called corticosteroids before your first 3 infusions of a treatment course
- You may also be given other medicines before or after the infusion to try to reduce your chances of having these reactions or to treat them if they happen. If something feels out of the ordinary, contact your healthcare provider right away

## WHY SHOULD I BE CONCERNED ABOUT POTENTIAL FOOD SOURCES OF LISTERIA?

People who receive LEMTRADA have an increased chance of getting an infection caused by the bacteria, listeria, which can lead to significant complications or death. Avoid foods that may be a source for listeria (for example, deli meat, unpasteurized milk and cheese products, soft cheeses, or undercooked meat, seafood or poultry) or make sure that the food you eat which may contain listeria is heated well if you receive treatment with LEMTRADA.

Please see Important Safety Information on pages 11-13 and full [Prescribing Information/Medication Guide](#), including serious side effects.

## WHAT SHOULD I BRING TO MY INFUSION CENTER?

It is recommended that you bring a sweatshirt or a blanket and wear comfortable clothes. You should pack something to entertain yourself during your treatment.

- Additional tips to help you pass the time during treatment can be found on page 4
- Be sure to call the infusion center to be certain what food, drinks, and entertainment are permitted and ask questions like whether water will be available for refillable bottles

## IS IT OKAY IF I GET PREGNANT WHILE TAKING LEMTRADA?

Before receiving LEMTRADA, tell your healthcare team if you are pregnant or plan to become pregnant as it can harm your unborn baby.

- Use effective birth control to avoid pregnancy during a treatment round
- Continue to use birth control for at least 4 months after a treatment round

## WHERE CAN I FIND ADDITIONAL SUPPORT?

An *MS One to One*<sup>®</sup> nurse is available 24/7\* to provide support if you're feeling nervous or have questions before, during, and after treatment.



## REMINDER

Be confident that you are ready to take this next step in your treatment journey.

\*As a member of *MS One to One*, you'll have access to an on-duty nurse 24/7. Regular *MS One to One* call center hours are Mon-Fri, 8:30 am-8:00 pm ET.

# IMPORTANT SAFETY INFORMATION

## LEMTRADA can cause serious side effects including:

**Serious autoimmune problems:** Some people receiving LEMTRADA develop a condition where the immune cells in your body attack other cells or organs in the body (autoimmunity), which can be serious and may cause death. Serious autoimmune problems may include:

- Immune thrombocytopenic purpura (ITP), a condition of reduced platelet counts in your blood that can cause severe bleeding that may cause life-threatening problems. Call your healthcare provider right away if you have any of the following symptoms: easy bruising; bleeding from a cut that is hard to stop; coughing up blood; heavier menstrual periods than normal; bleeding from your gums or nose that is new or takes longer than usual to stop; small, scattered spots on your skin that are red, pink, or purple
- Kidney problems called anti-glomerular basement membrane disease, which, if not treated, can lead to severe kidney damage, kidney failure that needs dialysis, a kidney transplant, or death. Call your healthcare provider right away if you have any of the following symptoms: swelling of your legs or feet; blood in the urine (red or tea-colored urine); decrease in urine; fatigue; coughing up blood

It is important for you to have blood and urine tests before you receive, while you are receiving and every month for 4 years or longer, after you receive your last LEMTRADA infusion.

**Serious infusion reactions:** LEMTRADA can cause serious infusion reactions that may cause death. Serious infusion reactions may happen while you receive, or up to 24 hours or longer after you receive LEMTRADA.

- You will receive your infusion at a healthcare facility with equipment and staff trained to manage infusion reactions, including serious allergic reactions, and urgent heart or breathing problems. You will be watched while you receive, and for 2 hours or longer after you receive, LEMTRADA. If a serious infusion reaction happens while you are receiving LEMTRADA, your infusion may be stopped.

Please see Important Safety Information on pages 11-13 and full [Prescribing Information/Medication Guide](#), including serious side effects.

Tell your healthcare provider right away if you have any of the following symptoms of a serious infusion reaction during the infusion, and after you have left the healthcare facility:

- swelling in your mouth or throat
- fast, slow, or irregular heartbeat
- trouble breathing
- chest pain
- weakness
- rash

To lower your chances of getting a serious infusion reaction, your healthcare provider will give you a medicine called corticosteroids before your first 3 infusions of a treatment course. You may also be given other medicines before or after the infusion to try to reduce your chances of having these reactions or to treat them if they happen.

**Stroke and tears in your arteries that supply blood to your brain (carotid and vertebral arteries):** Some people have had serious and sometimes deadly strokes and tears in their carotid or vertebral arteries within 3 days of receiving LEMTRADA. Get help right away if you have any of the following symptoms that may be signs of a stroke or tears in your carotid or vertebral arteries:

- drooping of parts of your face
- difficulty with speech
- weakness on one side
- neck pain
- sudden severe headache

**Certain cancers:** Receiving LEMTRADA may increase your chance of getting some kinds of cancers, including thyroid cancer, skin cancer (melanoma), and blood cancers called lymphoproliferative disorders and lymphoma. Call your healthcare provider if you have the following symptoms that may be a sign of thyroid cancer:

- new lump
- changes that do not go away
- swelling in your neck
- trouble swallowing or breathing
- pain in front of neck
- cough that is not caused by a cold
- hoarseness or other voice

Have your skin checked before you start receiving LEMTRADA and each year while you are receiving treatment to monitor for symptoms of skin cancer.

# IMPORTANT SAFETY INFORMATION (CONTINUED)

**Because of risks of autoimmunity, infusion reactions, and some kinds of cancers, LEMTRADA is only available through a restricted program called the LEMTRADA Risk Evaluation and Mitigation Strategy (REMS) Program.**

**Do not receive LEMTRADA if you** are infected with human immunodeficiency virus (HIV).

**Thyroid problems:** Some patients taking LEMTRADA may get an overactive thyroid (hyperthyroidism) or an underactive thyroid (hypothyroidism). Call your healthcare provider if you have any of these symptoms:

- excessive sweating
- unexplained weight gain
- unexplained weight loss
- feeling cold
- fast heartbeat
- worsening tiredness
- eye swelling
- constipation
- nervousness

**Low blood counts (cytopenias):** LEMTRADA may cause a decrease in some types of blood cells. Some people with these low blood counts have increased infections. Call your doctor right away if you have symptoms of cytopenias such as:

- weakness
- dark urine
- chest pain
- fast heartbeat
- yellowing of the skin or whites of the eyes (jaundice)

**Inflammation of the liver:** Call your healthcare provider right away if you have symptoms such as unexplained nausea, stomach pain, tiredness, loss of appetite, yellowing of skin or whites of eyes, or bleeding or bruising more easily than normal.

**Hemophagocytic lymphohistiocytosis:** LEMTRADA may increase the risk of overactivity of the immune system that can be fatal if not diagnosed and treated early. If you experience symptoms such as fever, swollen glands, or skin rash, contact your healthcare provider right away.

**Please see Important Safety Information on pages 11-13 and full [Prescribing Information/Medication Guide](#), including serious side effects.**

**Serious infections:** LEMTRADA may cause you to have a serious infection while you receive and after receiving a course of treatment. Serious infections may include:

- **listeria.** People who receive LEMTRADA have an increased chance of getting a bacterial infection called listeria, which can lead to significant complications or death. Avoid foods that may be a source of listeria or make sure foods are heated well.
- **herpes viral infections.** Some people taking LEMTRADA have an increased chance of getting herpes viral infections. Take medicines as prescribed by your healthcare provider to reduce your chances of getting these infections.
- **tuberculosis.** Your healthcare provider should check you for tuberculosis before you receive LEMTRADA.
- **hepatitis.** People who are at high risk of, or are carriers of, hepatitis B (HBV) or hepatitis C (HCV) may be at risk of irreversible liver damage.

These are not all the possible infections that could happen while on LEMTRADA. Call your healthcare provider right away if you have symptoms of a serious infection such as fever or swollen glands. Talk to your healthcare provider before you get vaccinations after receiving LEMTRADA. Certain vaccinations may increase your chances of getting infections.

**Progressive multifocal leukoencephalopathy (PML):** A rare brain infection that usually leads to death or severe disability has been reported with LEMTRADA. Symptoms of PML get worse over days to weeks. It is important that you call your doctor right away if you have any new or worsening medical problems that have lasted several days, including problems with:

- thinking
- eyesight
- strength
- balance
- weakness on 1 side of your body
- using your arms or legs

# IMPORTANT SAFETY INFORMATION (CONTINUED)

**Inflammation of the gallbladder without gallstones (acalculous cholecystitis):** LEMTRADA may increase your chance of getting inflammation of the gallbladder without gallstones, a serious medical condition that can be life-threatening. Call your healthcare provider right away if you have any of the following symptoms:

- stomach pain or discomfort
- fever
- nausea or vomiting

**Swelling of lung tissue (pneumonitis):** Some people have had swelling of the lung tissue while receiving LEMTRADA. Call your healthcare provider right away if you have the following symptoms:

- shortness of breath
- chest pain or tightness
- cough
- coughing up blood
- wheezing

**Before receiving LEMTRADA, tell your healthcare provider if you:**

- have bleeding, thyroid, or kidney problems
- have a recent history of infection
- are taking a medicine called Campath® (alemtuzumab)
- have received a live vaccine in the past 6 weeks before receiving LEMTRADA or plan to receive any live vaccines. Ask your healthcare provider if you are not sure if your vaccine is a live vaccine
- are pregnant or plan to become pregnant. LEMTRADA may harm your unborn baby. You should use birth control while receiving LEMTRADA and for 4 months after your course of treatment
- are breastfeeding or plan to breastfeed. You and your healthcare provider should decide if you should receive LEMTRADA or breastfeed.

**Tell your healthcare provider about all the medicines you take,** including prescription and over-the-counter medicines, vitamins, and herbal supplements. LEMTRADA and other medicines may affect each other, causing side effects. Especially tell your healthcare provider if you take medicines that increase your chance of getting infections, including medicines used to treat cancer or to control your immune system.

**The most common side effects of LEMTRADA include:**

- rash
- upper respiratory infection
- diarrhea
- headache
- sinus infection
- thyroid problems
- herpes viral infection
- mouth pain or sore throat
- fever
- hives
- tingling sensation
- swelling of your nose and throat
- itching
- dizziness
- nausea
- fungal infection
- stomach pain
- urinary tract infection
- joint pain
- sudden redness in face, neck, or chest
- feeling tired
- pain in your arms or legs
- vomiting
- trouble sleeping
- back pain

Tell your healthcare provider if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of LEMTRADA.

Please see Important Safety Information on pages 11-13 and full [Prescribing Information/Medication Guide](#), including serious side effects.

FROM BEFORE TREATMENT TO MONITORING,  
**YOU'RE ON YOUR WAY!**

If you have any questions,  
reach out to your *MS One to One*<sup>®</sup> nurse

Please see Important Safety Information on pages 11-13 and full  
[Prescribing Information/Medication Guide](#), including serious side effects.

SANOFI GENZYME 

© 2020 Genzyme Corporation. All rights reserved. LEMTRADA, *MS One to One*,  
Sanofi and Genzyme registered in U.S. Patent and Trademark Office.  
MAT-US-2008398 06/20

  
**LEMTRADA**<sup>®</sup>  
alemtuzumab<sup>12mg</sup>  
iv